DECEMBER 198

....



NUMBERS 18

# VINGNANAM JOURNAL OF SCIENCE

Published by

THE SCIENCE FACULTY
JNIVERSITY OF JAFFNA
SRI LANKA

# VINGNANAM JOURNAL OF SCIENCE

Editorial Board: G. F. RAJENDRAM (Editor), A. P. ARUDCHANDRAN, K. CHITRAVADIVELU, K. KUNARATNAM, R. MAGESWARAN, V. THARNIARATNAM (Associate Editors).

VINGNANAM – JOURNAL OF SCIENCE is a semi-annual journal devoted to the publication of original research papers and review articles in all fields of science and technology. Authors are entirely responsible for statements and opinion expressed.

The abbreviated title of this journal shall be Vingaanam J. Sci.

#### SUBSCRIPTIONS:

Annual subscription: Foreign \$ 20.00; Local Rs. 40.00 Single number: Foreign \$ 10.00; Local Rs. 20.00 Subscription includes postage.

Cheques should be drawn in favour of Bursar, University of Jaffna.

#### MANUSCRIPTS:

Manuscripts and related correspondence should be sent to Prof. G.F.RAJENDRAM, EDITOR, VINGNANAM — JOURNAL OF SCIENCE, Department of Zooleg, a University of Jaffan, Thirmselvely, Jaffan, Sri Lanka, Authors are urged to as "Intimactions to Contributors" printed at the end of this issue before preparing the manuscript.

#### COPYRIGHT

All rights are strictly reserved. No part of this journal may be copied, translated, photographed, broadcast, toler/sed or otherwise disseminated without the prior written permission of the Oean, Faculty of Science, University of Jaffan, Sri Lanke,

PUBLISHED BY THE

SCIENCE FACULTY OF THE UNIVERSITY OF JAFFNA
THIRUNELVELY, JAFFNA,

SRI LANKA.

DECEMBER 1989



VOLUME

NUMBERS 1 & 2



# VINGNANAM

JOURNAL OF SCIENCE

Published by
THE SCIENCE FACULTY

UNIVERSITY OF JAFFNA SRI LANKA

VOLUME 4



NUMBERS 1 & 2

# VINGNANAM

Publication of volume 4 (nos. 1 & 2) of Vingnanam has been delayed, once again due to circumstances beyond our control. The saddest among these is the untimely demise of the former Chief Editor S. Kandiah who had served in that canacity since the birth of Vingnanam.

In volume 4 we publish our first article in physics: "Diffusion of Hydrogen in Metals" by K. Kandasamy.

Two papers deal with diseases of local crops: the paper by R. V. S. Sundaresan and Pathmasany Varatharajah with the mosaic disease of papaya and the paper by Jayadovi Kailayapillai, A. Sivapalan and K. Theivendirarajah with the powdery mildew disease of sesame.

G. F. Rajendram with Nirmala R. Antony and Robini Nagaliagam bilitake the first study of the mosquito fauna in Jaffas periatula. Another Paper by G. F. Rajendram reports on goographic geotels variation among Populations of a green rice leafhopper from India, Sri Lanka and the Philippines.



CEL PUBLISH

MATERIAL TO A CO.

A front Million on the control of the College A ever for the applicability of the property of the control of th

model of three of these builts was to the factors of the following of the common of th

Contact has to come a major of medical tracolors of magnetic tractions and the part of the section of the come of the contact of the section of the sectio

#### VINGNANAM

# JOURNAL OF SCIENCE

Volume	4		December	1989	

Diffusion of Hydrogen in Metals. K. KANDASAMY ...

Instructions to Contributors

Volume 4

Numbers 1 & 2

14

. . .

# CONTENTS

A Study of the Free Amino Acids present in Leaves of Papaya (Carka	
papaya L.) varieties infected with Papaya Mosaic Disease.  R. V. S. SUNDARESAN and PATHMASANY VARATHARAJAH	
Biology of Peridomestic Mosquito Species of Jaffna Peninsula.	
G. F. RAJENDRAM, NIRMALA R. ANTONY and	

ROHINI NAGALINGAM	
Gel Electrophoresis of Enzymes of Nephotettin tera: Cicadellidae) populations from India,	Sri Lanka and the
Philippines. G. F. RAJENDRAM	y described to the

of Indiana I	). JEYADEVI KAILAYAPILLAI, K. THEIVENDIRARAJAH	2
a. Sitarana		4

A. SI	VAFALAIN AND IL		
41		 	
Abstracts in	Tamil		

#### PALAN

GZ - 19 3 3 2 2 19

	CONTRACTOR	
of Jaffna e	×	

character of regardal to adjust

The state of the s

And the state of the second of

The side interpretation of Engineer of Mydiates recovered some for title different purchases from the side to the same on Principles of Englishments.

The second sector is a second second

lead to see the

#### DIFFUSION OF HYDROGEN IN METALS

#### K. KANDASAMY

(Department of Physics, University of Jaffna, Sri Lanks.)

#### Vingnanam — Journal of Science 4: 1 — 7 (1989)

ABSTRACT: The Chen Min Li formalism (Chen Min Li 1939) for the flor of mobile solute in a stressed solid is used to derive an equation for diffication flux of hydrogen in matrix. The predictions from this equation see compand with the apportmental results exported by Livet's r.d. (1937). The agreement is additationly.

## entities the area representation are at the areas the X and the

Inclusion of Hydrogen into the lattice of many metals causes expansion of their lattice (Kandasamy 1980), which in turn creates an elastic field in the metal lattice. The created elastic field is not limited to the sites of inclusion but is extended over a long range in the metal. The elastic field in a metal which contains hydrogen is capable of inducing diffusion of hydrogen (Gorsky 1935). The diffusion processes in metals, which exhibit lattice expansion on the inclusion of hydrogen, therefore, cannot be governed by Fick's law. Fick's law assumes that the concentration gradient is the only physical parameter which contributes to the diffusion of hydrogen in metals. From thermodynamic principles, Chen Min Li developed an expression for the flux of mobile solutes in a stressed solid (Chen Min Li 1978). Using network model Larche and Cahn (1982) derived an expression for the diffusion flux of bydrogen in metals. It was later amended by the author (Kandasamy 1988). More recently non-Fickian nature of hydrogen diffusion in metals, particularly in talladium based alloys (Pd Pt 19) was experimentally demonstrated by Lewis and his coworkers (Lewis et al. 1987). In this communication an equation for the diffusion flux is obtained from the Chen Min Li expression and this equation is solved using computers to explain the results of Lewis and his coworkers (1987),

#### Theory

In a solid with inhomogeneous distribution of hydrogen the hydrogen diffusion is not only governed by the concentration gradient but also by the gradient of the stress which is induced by the inclusion of hydrogen the the metal. According to Chen Min Li (1978) the diffusion flux of hydrogen

in the x direction, Jx, when a metal membrane, which initially contained a uniform distribution of bydrogen concentration, Co, absorbs further hydrogen into one of its faces parallel to the YZ plane, is given by

$$J_{X} = -D\frac{dc}{dx} + \frac{Co}{R}\frac{\overrightarrow{V}_{H}}{T}\frac{D}{dx} \qquad (1)$$

where D is the diffusion coefficient of hydrogen in the metal,  $\overline{V}_{H}$  is the partial molar volume of hydrogen in the metal,  $\frac{dc}{dx}$  is the concentration gradient

dd is the stress gradient and the other symbols have their usual meaning. The far induced stress 6 depends on the geometry of the metal sample in addition to, we shall consider a thin shab sample of thickness L (Fig. 1) containing a uniform distribution a dylorgen of concentration Ca. If one of its faces, say the face at X=O and parallel to the YZ place, is exposed to a high concentration Ca of bydrogen, the absorption of hydrogen distribution they have the mean state of the distribution of hydrogen distribution to the YZ place, it exposed to a high concentration of hydrogen disturbs the initial uniform distribution of hydrogen and incluses stress in the metal lattice.




Fig. 1: Variation of hydrogen concentration (C) along the thickness of a thin slab (thickness of slab=1). Co is the initial concentration (distributed uniformly) of hydrogen in the slab. Cs is the concentration of the hydrogen atmosphere into which the absorbing face (face at X=0) of the slab is exposed.

The new distribution of hydrogen concentration is represented by  $\frac{C-Cs}{Cs-Co} = 1-2\sum_{\substack{n=0\\n=0}} \binom{1}{n} \rho_0 \exp \begin{bmatrix} \binom{-\rho^2}{n} Dt \\ \frac{1}{L^2} \end{bmatrix} \cos \rho \begin{bmatrix} X \\ \overline{L} - 1 \end{bmatrix}$ 

$$= \frac{\Delta C}{\Delta C_8} \text{ (say)},$$
where  $\frac{\rho}{n} = (2n+1)\pi/2$ 

The three components of induced stress are given by

The three components of induced stress are given by  $d_{XX} = 0$ 

$$\begin{split} & \stackrel{\circ}{\text{yy}} = \stackrel{\circ}{\text{yz}} \\ & \stackrel{\circ}{\text{y}} = -\frac{-V_N}{y(1-y)} \left\{ \triangle C - \frac{1}{L_0} \int_0^z \triangle C \, dx - \frac{12}{L^2} \left\{ X - \frac{L}{2} \right\} \int_0^z \triangle C \, \left\{ X - \frac{L}{2} \right\} dx \right\} \end{split}$$

(2)

where E is the Young's modulus, and  $\nu$  is the Poisson's ratio of the sample and  $\Delta C$  is the change in hydrogen concentration due to absorption. The above equation is obtained from the Chen Min Li's equation with the imposition of boundary conditions used in this study.

On substitution for  $\frac{d_0}{dx}$  in equation (1) for the flux of hydrogen, we will get

$$Jx = -\left\{1 + \frac{2Co\bar{V}^2\pi E}{3(1-v)RT}\right\} D \frac{dc}{dx} + \frac{8CoE\bar{V}^2\pi D}{(1-v)RTL^2} \int_{0}^{L} \Delta C\left[X - \frac{L}{2}\right] dx$$
(4)

The form of the above equation is similar to the amended equation of Larche and Cahm (Kandssamy 1988). The first term in the equation represents the modification in the effect of concentration gradient in diffusion due to the presence of hydrogen is the metal lastice. The second term represents the effect of induced stress due to the inclusion of hydrogen in the metal lattice.

### Results and Discussion

We shall consider an experimental situation like the one conducted by Lewis et al. (1987) in which the hydrogen atoms diffusing through a tube having a thin wall of thickness L and area A are collected in a volume V. Por this case, the pressure change P, inside the volume is given by

Per this case, the pressure change 
$$P$$
, usade the  $P = \frac{RTA}{V} \int_{X=L}^{d} dt$  (5)

The thickness of the tube wall is very small compared to the diameter of the tube, and therefore it can be considered as a thin slab.

Vingnanam - Journal of Science Vol. 4, Nos. 1 & 2

The flux diffusing out of the slab is  $J_{X=L}$  which is given by  $J_{Y=1} = - \left\{ 1 + \frac{2 \text{ Co } \vec{\nabla}^2 \text{ is E}}{3(1-\epsilon) \text{ RT}} \right\} \quad \text{D} \quad \text{dis} \quad X \to L \quad \frac{8 \cdot \vec{\nabla}^2 \text{ in EDCo} \triangle \text{CsB}}{1 - \epsilon}$ 

Where 
$$B = \sum_{p} \frac{(-1)}{p^2} \begin{cases} p \sin p + 2 - 2 \cos p \\ n & n \end{cases} \exp \begin{cases} -p^2 D_1 \\ \frac{1}{L^2} \end{cases}$$

For metals like palladium  $\nabla u = 1.65 \times 10^{-6} \text{m}^3 \text{ mole}^{-1} \frac{E}{1-y} = 10^{11} \text{ Nm}^{-2}$ and  $C=10^{20}$ . where n is the hydrogen atoms to metal atoms ratio. At room temperature  $RT=5 \times 10^{3} \text{ J}$  mole-<sup>1</sup> On substituting these data





Fig. 2: Variation with time (t) of the pressure (P) inside the volume V (into which the diffused hydrogen atoms are being collected) computed using equation (8) with △ns=0.05 and different values for no



conditions: membrane equilibrated from both sides with a hydrogen pressure of 40 Torry pressure increase in the bulb (indicated by arrew) to 625 Torr: temperature, 26+9-2° C; breakthrough time, 12.5 min. Fig. 3: Experimental pressure (P) variation inside the volume V (into

as a function of time. Initial

as a function of time. Initial conditions: membrane equilibrated from both sides with hydrogen pressure of 605 Torr; pressure increase in the bulb (indicated by arrow) to 815 Torr: temperature, 26±0.2° C; breakthrough time, 19-3 min.

which the diffused hydrogen atoms are being collected) with time (t). (Lewis et al. 1987)

The line 1 in Fig. 2 shows the variation of P inside the volume V when the absorbing surface of the tube, which has no initial hydrogen, is exposed to a hydrogen atmosphere corresponding to  $n_8$  = 0.05. As has been expected the variation of pressure reflects the FICKIAN nature of hydrogen diffusion. Lines 2 and 3 in the same figure show the variation of pressure in the volume V when the absorbing surface of the same tube, but with au initial non zero uniform hydrogen concentration, is exposed to the hydrogen atmosphere of same pressure difference. For lines 2 and 3 the initial hydrogen concentration in the tube wall corresponds to no=0.01 and 0.02

respectively. The initial drop in pressure indicates that immediately after the exposure of the absorbing face of tube wall to the hydrogen atmosphere, the hydrogen atmosphere was the hydrogen atmosphere with the hydrogen atmosphere was the hydrogen atmosphere with the hydrogen atmosphere was the hydrogen atmosphere with the hydrogen atmosphere with the hydrogen atmosphere with inference at the hydrogen atmosphere with increasing the of the hydrogen atmosphere with the hydrogen a

The non-Fickian diffusion is the consequence of induced steas in the metal lattice by the presence of bydrogen atoms in them. To illustrate the nature of effect produced by the induced steas in the hydrogen diffusion, the pressure variation does to contribution coming from the second term alone in the flux equation (8) is plotted in Fig. 4. The on-Fickian effect increases with increasing ACs or (2, ns). This is clearly demonstrated in Fig. 4. Further, the plots in the Fig. 4 indicate that the effect of stress is not only limited to initial stages but extends to the entire experimental range. Therefore all the data obtained at the steady state by Lewis et al. (1987) include the effect of none-fixing diffusion follydrogen. Values deduced from this data for quantities such as coefficient of diffusion, solubility exp. using Fick's law are therefore erroneous.




Fig. 4: Variation with time (t) of the pressure (P) inside the volume V (into which the diffused bydrogen atoms are being collected) computed using only the second term of equation (8) with different values for Δ n.

#### Conclusion ·

This study gives a simple theoretical analysis of the data obtained by the data obtained by the data obtained by a simple data obtained by the data obtained by the data of th

#### Acknowledgement

My thanks are due to Dr. F. A. Lewis, Queen's University, Belfast, Northern Ireland, for his valuable advice and to Mr. S. Kannan, Computer Unit, Université of Jaffina, Sri Lanka, for assistance in computing.

#### Reference

- Chen Min Li, J. (1978) Physical chemistry of some microstructural phenomena, Metal. Trans 9A, 1353-78
  Gorsky W. S. (1935) Hydrogen diffusion in solids. Phys. Z. Soviet Union 8.
- 457.

  Kandasamy, K. (1980) Interaction of hydrogen and oxygen with titanium films.
- Ph. D. thesis, University of Keele, England.

  Kandasamy, K. (1988) Hydrogen concentration distribution in elastic membranes
- during diffusion of hydrogen. Scripta Metallurgica 22, 479 481. Larche, F. C. & Cahn, J. W. (1982) The effect of self stress on diffusion in
  - solids (Acta Metals) 30, 1835-45.
- Lewis, F. A. Baranowski, B & Kandasamy, K. (1987) Uphill diffusion effects induced by self stresses during hydrogen diffusion through metallic membranes. Journal of the Less Common Metal 134, L27-31.

# A STUDY OF THE FREE AMINO ACIDS PRESENT IN THE LEAVES OF PAPAYA (CARICA PAPAYA L.) VARIETIES INFECTED WITH PAPAYA MOSAIC DISEASE

# R. V. S. SUNDARESAN & PATHMASANY VARATHARAJAHI (Department of Botany, University of Joffes, Srl Lanks)

Vingnanam — Journal of Science 4: 8 — 13 (1989)

ABSTRACT: The fine minos acids present in leaves of Carcaphopys. L. C. V. Colembates 2, D. Plas, L. Cook, Sido Hessand subnoyus Indiana, both healthy and affected by the papies mostel disease were determined by two dimensional paper chromotography, since subjecting the leaf extracts to loss subnergo chromotography. The amino acids allosine as—more bodyire acid, aprillers, president acid, cystelic acid, glotassic acid, cylorus, hestilinis, isolaroctic acid, prised acid, glotassic acid, cylorus, fine colorus, cytteliosections, brain, methodorus, phanyl adams, partin, placesic, cytteliosections, brain, methodorus, phanyl adams, partin, placesic, cytteliosections, prise, methodorus, phanyl adams, partin, placesic, cytteliosections, partin, methodorus, partin, partin, partin, partin, partin, partin, settledis, but in verpley, concentrations in different varieties, and the healthy and infected varieties and the variation in their concentrations.

#### Introduction

Vital Infection of plasts results in the protein and nucleic acid synthesizing membrary of the bost cells bring intense over by the virus. As a consequence, the protein capsid and the nucleic acid of the virus are synthesized in preference to the proteins and nucleic acids of the host. In addition other proteins as pacified by the boat and which are envolved in inducing resistance against which are conditions of the protein as pacified by the boat and which are tenvolved in inducing resistance against which are conditions to the protein and are likely to be synthesized in significant quantities in the cell. That was decided to investigate whether there were any differences in the composition and concentration of amino acids from leaf extracts of papays plantic systemically infected with passing mosaic disease compared with healthy systemically infected with passing mosaic disease compared with healthy and the protein of the protein acids to difference in a supposition of the protein constraints of acids to difference in the acceptability or however.

<sup>1</sup> Present address: School of Agriculture, Kundasale, Sri Lauka.

Previous investigati as undertakan with the same virus-host system had been carried out only in the variety Solo Hawaii and had revealed only (Rajapake, 1981). This observation solds, both in the healthy and infected leaves in order to obtain a more thorough and true picture of the anino acid composition in viral infection.

It has been recently reported that the causative organism of papaya mosaic disease resembles in many aspects the papaya ring spot virus (PRSV) and is suspected to be a strain of this virus (N. Shivanathan; personal communication).

#### Materials and Methods

### Varieties of Papaya under study

Five Papaya (Carine prayas L) varieties were selected for this study. These are C. V. Coimbaro et a. CPL value Shot Hawsit and a unknown Indian variety. Healthy Seeds of four of the varieties were obtained from the Central Apricultural Research Institute, Gannouwa, and the lood variety seeds were obtained from a farmer in Jaffan. Seedlings were nined in polythese baps. Five completely readounded design, a capriment was laid out according to the completely readounded design.

#### Source of virus and inoculation procedure

Virus Virus infected Solo Hawaii leaves obtained from Kandy district were used to inoculate all five varieties of papaya plants.

Preparation of inocolum. One gram portions of infected leaves, after deribbing, were thoroughly ground in phosphate buffer in a sterile mortar. The extract was filtered through a piece of muslin cloth and the filtrate made up to 10 ml with phosphate buffer. A pinch of Carborundum was mixed thoroughly with the filtrate as an abstaint.

Insculation of plants. Replicate plants were kept in a dark chamber 260 mbrs before incordation. The following afternoon the plants were removed 760 mbr chamber and the first pair of leaves were inoculated by probbing the insculum on the first super surface with the forelines. The incovated leaves were traved with distillation of the fact of the first proper surface with the first probability of the fact of the fact surface. Controls were also maintained all the plants were maintained in the stead proof glists-house.

#### Amino Acid Analysis

Preparation of the plant extract. The preparation of the plant extract and subsequent ion exchange chromatography of amino acids was done according to Jeyaratnam (1986). Ten grams of fresh leaves of both healthy and infected plants, two weeks after inoculation, were separately ground in a V-1

10

blender with 50 ml of 90% ethanol. This homogenate was centrifuged at  $10 \times 1000$  r.p.m. for 30 min. The pellet was discarded and the supernate ensporated to dryness in a crucible. The residence was disorded in 25 ml of the period of the p

Preparation of ion exchange column and collection of free name ratio. A glass column of 1 cm diameter was packed with washed calicon exhaps resin (Dower 59) to a helgit of 20 cm. Before use the resin column was converted to the 1<sup>st</sup> form by passing 25 ml of 28 HG through it followed by distilled water to wash away the unabsorbed chloride ions. The even until the washings were free of chloride ions.

The aqueous plant extract was then allowed to run dowly thoughts of this column. The flow rate was regulated to bring about efficient planting of the amino scids. This was followed by washing of the column with distilled water. The aborebed amino acids were then eluted with 25 m if 3N NH,OH. Again 15 m if 3 N NH,OH was added to cluste the remaining anison acids. The cluster was collected in a cleen glass beaker and eviporated amino acids. The cluster was collected in a cleen glass beaker and eviporated acids of the contract of the contract of the contract of the cluster was contracted the free amino acids. Aliquots of this were used for two dimensional paper (chromstographs)

Separation of amino acods by paper chromatography. Descending two dimensional paper chromatography was carried out using Wataman so I chromatography paper, as described by Elvidge and Sammes (1966); in 10<sup>3</sup> luliquoid with the extracts were spotted and the paper was developed with Batan—I-ol's acute said water (4:1:1 railo) in the first direction and with plane's cortic said water (4:1:1 railo) in the first direction and with plane's railor of the said water (4:1:1 railo) in the first direction acods were leaved by spraying with 0.5% inishydin in butan—I-ol and warmed in the overallor SPF. The amino acids were later identified by comparison with a stand-chromatogram and the concentration determined by visual observation. The proceedings was repeated thrite for both healthy and interest leaves of each of the contract of

#### Results and Discussion

of both healthy and infected leaves are outlined in Table 1. Arginine was not detected either in healthy or infected leaves of the varieties Coimbatore - 2 and Solo Hawaii and in the infected leaves of the varieties CP 124 and local variety, Table 1.

Free amino acids and their relative concentrations in leaves of different Papaya varieties

		163	ives of	dille	rent Pa	ipaya	varietie	s		
Amino acid	Coimbat Healthy		CP Healthy		Lo Healthy	ral Infec.	Solo H Healthy	nwali nfec.	Unknows	Indian Infec.
1) Alanine	М	M	w	s	s ·	М	М	s	s	М
<ol> <li>amino butyric acid</li> </ol>	т	T	М	м	М	М	w	M	s	- S

3) Arginine M 4) Aspartic acid M

5) Cystelo Т acid 6) Glutamic S S S acid S M м М M 7) Glycine м м M w W

т T 3.6 8) Histidine т M м S 9) Isoleucine M S M т M 10) Leucine M s w т т w M 11) Lysine s S s s 12) Methionine M: S

13) Phenyl м M т W alanine M S S S S s s 14) Serine w w

м w M M м M w w M T т 16) Tyrosine T s s S 17) Valine s S т Ŧ

15) Threonine

M - Moderate, W - Weak, T - Trace

Scoring scale: S - Strong.

12

The absence of this amino acid in Coimbatore -2 and Solo Hawaii may be a varietal character, whereas its absence in the infected leaves of CP 124 and local variety too, may indicate the relatively less important role arginine plays in the synthesis of the capsid.

The concentrations of evsteic acid and lysine are less in i fected leaves of all five varieties compared with the corresponding healthy leaves. This sorees with the observations made by Rajapakse (1981). The reduction in the concentrations of different amino acids in only the infected leaves of different varieties may be due to the heavy demand of these amino acids for the synthesis of the capsid.

In the variey Coimbatore- 2, histidine, isoleucine, leucine, lysine, methionine and nhenyl alanine were present in higher concentrations in the infected leaves than in healthy leaves. It is possible that new proteins may be synthesized from these amino acids in infected plants which may be present as a factor in resisting or tolerating virus infection (Kimmins, 1969). Observations indicate that the presence of high concentrations of amino acids is associated with low infectivity of the causative virus (Varatharaiah, 1989). On the other hand there was a reduction in the concentration of six amino acids in the infected leaves of the variety Solo Hawaii, four in the infected unknown variety, three in the infected local variety and one in the infected CP124 compared with the corresponding healthy leaves. It is possible that the reduction in concentration of many amino acids in the infected leaves of the above varieties may point to a higher incidence of infection and greater susceptibility of these papya varieties to mosaic disease.

#### Acknowledgement

This study forms part of the M, Sc. thesis of the junior author who was granted study leave by the School of Agriculture Kundasale.

#### References

Elvidge, J. A. and Sammes, P. G. (1966) Paper chromatography pp. 40-46. In: A Course of Modern Techniques of Organic Chemistry Butterworth, United Kingdom

Jeyaratnam, M. (1986) The Chemistry and Biochemistry of Palmyrah Products. M. Phil thesis, University of Jaffna, Sri Lanka,

Rajapakse, R. H. S. (1981) Biochensical analysis of healthy and mosaic virus infected papaya leaver. Madras Agricultural Journal, 68 (7), 458 - 440.

Varatharajah, P. (1989) Susceptibilty and physiological studies on Papays (Carica papays L.) varieties to Papaw Mosaic disease, M. Sc. Thesis University of Jaffna, Sri Lanka.



received for publication June 196



# BIOLOGY OF PERIDOMESTIC MOSQUITO SPECIES OF JAFFNA PENINSULA

# G. F. RAJENDRAM, NIRMALA R. ANTONY AND ROHINI NAGALINGAM

(Department of Zoology, University of Jaffns, Sri Lanks)

Vingnanam — Journal of Science 4: 14-19 (1989)

ABSTRACT: The biology of three professestic meaguine secretics from Jeffern periodes was straight Advantage Linneaus, Armigeria de-lamit Sewaria and Culice policypic Linneaus, 45 days and pous 67 days. The total developmental profess was professed for the few homestall pages of all and tentum was a gap 24 days, and the companies of the few homestall pages of all the few homestall pages of the separation of the few homestall pages of the few ho

#### Interoduction

Studies on monguines of Sci Lanka, including the recent Southeast Asis Monguite Project and Medical Entomology project of the Smithonian Institut, Washington D. C., liave been confined to the southern and central regions of Sci Goren 1903, Corter 1903, Kombelin 1981, Ameriaghe 1981, Studies and Corten India openies and India peniestal include a recent survey of periforments most in northern Jaffra peninsula include a recent survey of periforments benchmark of the Corten 1981, American 1981, The president of periforments in the India of Action and April 1981, American 1981, The president periforment of the India of Periforments in mosquito species of Maintenance and Periforments and Cortenance and Corte

#### Materials and Methods

Live mosquitoes of the three species were collected in individual glass vials and sorted out under a dissecting microscope. They were transferred to plastic sleeve cages 30×25×25 cm, evered with fine nylon mesh at the side and with a plastic door in front and a cloth sleeve fitted on one side. A

December 1989 Rajendram et al.: Biology of Mosquitoes of Jaffna

plastic container 7.5 cm diameter and 10 cm high with a rice plant was placed in the cage as well as a 200 ml beaker half filled with water. Observations were made daily. After the eggs hatched into larvae, powdered maldive fish at the rate of 12 gm/litre, was added daily as food.

The temperature in the laboratory registered a range of 26-32° C and relative humidity 40 to 92%.

### Results

The total developmental period of A. a-gypti from egg to adult stage averaged 12.9 days; the egg stage averaged 1.69 days; the larval stage 4.5 days; and the pupal stage 6.7 days (Table 1).

Table 1: Developmental period of A. argypti

Stage	No. of days	No. of eggs	Average duration in days
Egg	5.0 83-2 4 <sub>1</sub> 2 51 o 514	23	1.7
200	2	26	
	3	6	1 67±0-66
Larva	the state of the state of	20	
Laiva	4	10	
A section		10	45.00
	6	7	
	7	8	4.5 ±146
to re-	ide to K	2	
Pupa	3	18	
	5	3	
	100	10	
		18	
	9	4	6-7 ± 1-5

Total developmental period = 12.9 ± 3.62 days

	Stage	No. of days	No. of eggs	Average duration in days
averag days;	ged 14.0 days: and the pupa	the egg stage av I stage 4.04 days	of A. durhami from eraged 3.12 days; the (Table 2). period of A durham	larval stage 6.
		nanam - Journal o	of Science	Vol. 4, Nos. 1 &

	4	10	3·12±0·83
Larva	5	3	
	6	10	
	8		6·84 ± 1·18
Pupa	1	3	

4·04 ± 1·65

Total developmental period = 14.0 ± 2.66 days

The number of eggs per raft, laid in the laboratory, by C quinquefasciaus

averaged 112.8 (Table 3). Table 3: Frequency distribution of number of eggs per raft among

12 unfragmented egg-rafts of C. quinquefasciatus

No. of eggs No. of rafts Average no. of

eggs per raft

80 - 89 01

90 - 99

01

100 \_ 109

02

110 - 11904

120 - 129 02 130 - 139 112·8±13·7

02

The total developmental period of C. quiquefaccious from egg to adult stage averaged 14.3 days; the egg stage averaged 5.6 days; the larval stage 5.8 days; and the pupal stage 2.9 days (Table 4).

Table 4: Frequency distribution of minimum duration of various stages of 12 egg rafts C. quinquefascianus

Stage	No, of days	No. of raits	Average duration in days
Egg	2	2	
	3	4	
	4	1	
	7	2	
	8	1	
	12	1	
	13	1	5·6±1·9
Larva	.5 7 3	2	
		set i .	
	7	2 .	
	8	4	5-8±2-I
Pupa	1	1	
		. 3	
	3	4	
	4	4	29±09

Total developmental period = 14:3±3:8 days

#### Discussion

The total developmental period of A. segpri in the present study was 129 days, with the larval stage at 4.5 days. Framin' (1997) reported the minimum period of larval developments A. secrety at a constant temperature of the contract of the contract temperature of the contract temperature of the contract temperature of the contract of the contract temperature of th

Studies on the biology of Culer pripers forigent. Weid, I.C. quirt-prissing, in Sri Lanka were carried out by Chow and Thexassaysmi (1973), Tay reported that the duration of the period between egg and adult averaged 31-20 days; the egg stage was 1-2 days (average 15-6 days); the Laral stage was 1-2 days (average 16-86 days); the Laral continuation of the pupul stage, 1-4 days (average 19-8 days). They obtained similar results with egg-rafts kept in water obtained from earth pits instead of water obtained from a well. In Calcutta, Roy (1949, reported the period between egg and adult of C. quinterprissional was 3-10 days during January, St-F minimum and 81° F maximum (12-2-72°C). Pennervy (1929) reported, without mentioning the emperature, that the minimum duration of the egg stage was 24 hours, of the larval stages, 120 hours, and of the pupul stage, 48 hours.

The developmental period of the three species are related to the laboratory conditions of temperature 26-32° C-

# Acknowledgements

Grateful thanks are due to the following of the University of Jaffuz Prof. V. K. Ganesalingam, Dean, Faculty of Science, and Mrs. N. Selvarajah, Head, Department of Zoology, for facilities; Mr. S. Narendran for technical assistance and Mr. S. Sriskantharasan for field assistance,

Thanks are also due to Dr. P. S. Cranston and Mr. B. C. Townsend of the Entomology Department, British Museum (Natural History), London, for taxonomic assistance.

A portion of this study formed part of a M. Phil, thesis submitted by the second author to the University of Jaffna. This work was supported in part by grant No. RG/85/8/6 from the Natural Resources, Energy and Science Authority of Sri Lanka awarded to the first author.

#### References

Amerasinghe, F. P. (1982) Observations on the mosquitoes (Diptera: Culicidae) of Udawattakele Forest, Sri Lanka. J. Natn. Sci. Coun. Sri Lanka 10 (1), 81 – 97.

Antony N. R. and G. F. Rajendram (1987) Mosquito oviposition in bamboo trape in Jaffna Peninsula. Proc. Sri Lanka Assoc. Advmnt. Sci 43,137 (Abstract).

Carter, H. F. (1950) Ceylon mosquitoes: lists of species and names of mosquitoes recorded from Ceylon. Ceylon J. Sci. (B) 24 (2), 85-115.

- December 1989 Rajendiam et al.: Biology of Mosquitoes of Jaffina 19

  Chow, C. Y. and E. S. Thevasagayam. (1957) Bionomics and control of Culex pipiens futigans Wied. in Ceylon. Bull. Wild. Hith. Org. 6,
  - cutex pipens fungans Wied in Ceylon, Bull, Wld, Hlth, Org.
- Francis, E. (1907) Observations on the life cycle of Stegomyla calopus. Publ. Hlth. Rep. Wash. 22, 381 - 383.
- Green, E. E. (1901) Mosquitoes and malaria. Royal Botanic Gardens. Peradeniya, Circular Series, 1. No. 25, 345 –358.
- Peradeniya, Circular Series, 1. No. 25, 345 358. Krombein, K. V. (1981) The Smithsonian insect project in Sri Lanka. Spolia
- Zeylanica 35 (1 & 2), 119 134.

  Mitchel, E. G. (1907) Mosquito Life. G. Putnam's Sons, New York.
- Pomeroy, A. W. J. (1920) J. Roy. Army Med. Cps. 35, 44.
  Rajendram, G. F. and N. R. Antony (1986) Mosquito sampling in several
- Rajendram, G. F. and N. R. Antony (1986) Mosquito sampling in several locations in Jaffaa Peninsula. Proc. Sri Lanka Assoc. Advmnt. Sci. 42, 144 - 145 (Abstract).
  Rov. D. N. (1946) Entomology. Excelsior, Calcutta.

Received for publication November 1990

Al tribut Logica and De Nova Story of the Artist Control of the Ar

# STARCH GEL ELECTROPHORESIS OF ENZYMES OF NEPHOTETTIX VIRESCENS (HOMOPIERA: CICADELLIDAE) POPULATIONS FROM INDIA, SRI LANKA AND THE PHILIPPINES

#### G. F. RAJENDRAM 1

(School of Pure & Applied Biology, University of Wales, Cardiff CF1 3TL, United Kingdom)

Vingnanam - Journal of Science 4: 20-26 (1989)

ABSTRACT: The banding patterns of eight soluble enzymes. esterase (EST), glucose-6-phosphate dehydrogenase (G-6-PDH), hexokinase (HK), isocitrate dehydrogenase (IDH), lactute dehydrogenasa (LDH), malate dehydrogenesa (MDH), malic enzyme (ME) and phosphoglucomutase (PGM), were studied in Nethicettix virescens (Distant) populations collected from India, Sri Lanka and the Philippines and mainteined in class-houses, on rice variety TN1, at University of Wales, Cardiff, United Kingdom, Adult Insects of both sexes were analyzed, using starch gel electrophoresis, and the gel stained for the eight enzymes. The bands in seven enzymes agreed in number and relative mobility in the three populations. In esterase, though three zones of activity wore observed in all three populations. the relative mobilities of two bands in the Philippines population differed from N. viroscess of India and Sri Lanks, and could be used for discusstic separation of the Philippines population. Esterase - 1 bands appear to be sex-related since they were more Intense in the females. The regults some with previously reported divergence in the acoustical behaviour of N. virescens from Sri Lanks and the Philippines, inducing partial reproductive isolation between these populations.

#### Introduction

petity, has been made possible by electrophyretic analysis of enzymes (gestluml petity, has been made possible by electrophyretic analysis of enzymes (Berlothe 1979, Powell et al. 1980. Ayala 1983). One such group of innects studed are the green leafthoppers, Nephaterix species, which are common pets of lite rice crop in Southeast Asia, Australia and the Pacific Islands (100se 1986). These innects have grown in importance as major pets of fice due to lite failing to transmit virus diseases (Radole 1983.) Genetic differences between

<sup>1</sup> Present Address: Department of Zoologe, University of Joffins, Sri Lenks

21

December 1989 Rajendram: Enzymes of N virescens

Nephotettix species have been studied using star-h-gel electrophoresis. Ozaki (1969) reported on esterase variation in relation to organophosphate r. sistance (1989) reported on genetic in N. p. otettix cincticeps (Uhler). Rajendram (1987) reported on genetic in N. p of the control of the contro present paper reports on geographic genetic variation between Nephatestiv present pupos reported as a contract of native cultivars of rice varieties resistant to various pests (Khush 1979), Sri Lacka, an adjacent island country, also a similar source of resistance, and the Philippines, a distant country, devoid of native cultivars of rice with resistant genes.

#### Meterials and Methods

N. virescens was collected from rice fields in Cuttack, India, Kandy, Sri Ianka and Los Banos. Philippines and maintained on rice variety TN-1. in temperature controlled glass-houses at University of Wales, Cardiff, United Kingdom, N virecens was identified according to the characters of the vertex. the nature of the submarginal and marginal black bands on the vertex and the black marking of the anterior of the proportum, and the senitalia of the 4th instar nymphs (Ghauri 1971).

Electrophoretic analysis was carried out on a total of 80 adults, 40 males and 40 females, from each population for each enzyme system. Individuals from the different populations were run on the same gel to show that any mobility differences were not due to inconsistencies between individual gels. Starch gel was prepared using 11.25% Connaught hydrolyzed starch (lot no. 425-1) in a tris-borate EDTA (ethylenediaminetetra-acetic acid) buffer (pH 8-6). Details of sample preparation and electrophoretic technique are given by Rajendram (1987). An insert dipped in 1% solution of buffered (pH. 8.6) bromophenol blue served as a reference for calculating isozyme migration distance. The bromophenol blue migrated approximately 13 cm during running of the gel, which lasted 5 hours. After electrophoresis, the gel was sliced horizontally and the slices stained for the following enzymes; esterase (EST) (E.C.3.1.1). glucose-6-phosphate dehydrogenase (G-6-PDH), (E.C.111.49), hexokinase (HK) (E.C.2,7.1.1.), isocitrate dehydrogenase (IDH) E.C.1.1.142), lactate dehydrogenase (LDH) (E.C.1.1.27), malate dehydrogenase (MDH) (E.C.I.I.1.37), malic enzyme (ME) (E.C.1.1.40) and phosphoglucomutase (PGM) (E.C.2.7.5.1) according to the methods described by Shaw & Prasad (1970). Among the enzyme systems studied, the fastest band, the most anodal, was assigned a mobility of 1.0, and the other bands were assigned relative mobilities when compared to the distance travelled by the standard band. The slowest band was the most cathodal. In each enzyme system, the bands were numbered 1, 2 or 3 in order of decreasing relative mobility.

#### Results

# 22 Esterase

In esterase, three zones of activity were present in all three populations of N-Nexesset (FE). D. Est-1 comprised three weakly staining bands with replace mobilities. They were present, with virtually identical relative mobilities at 10, 00,7 and 0.94, in the males and feenales of all three population. These appear to represent a polymorphic locus with three allest, apparently monomeric in structure. But the hands were insufficiently stained for fellule scoring. All three hands in Est-1 stained more intensely fainting band and was found in both males and females with relative mobilities of 0.38 in the India and fri Lanta populations and 0.29 in the Philippines population. Comprised a band with equal intensity with a relative mobility of 0.030 in the India and fri Lanta populations and 0.19 in the Philippines population.




Fig. 1 Zymograms of esterase (EST) patterns in N virescens populations from (A) India, (B) Sri Lanka and (C) the Philippines.

#### Glucose-6-ph sphate dehydrogenase (G-6-PDH)

Two monomorphic bands were seen at relative mobilities of 0.20 and 0.14, identical in all three populations.

### Hexokinase (HK)

One zone of activity was present, with an apparently monomorphic hand, with a relative mobility of 0.47 in all three populations.

# Isocitric dehydrogenase (IDH)

Only one zone of activity was observed. The males possessed a fast electromorph, a single, sharp, well resolved band, with a relative mobility of 0.20 and the females a slightly slower and more diffuse band with a relative mobility of 0.14. The relative mobilities were virtually identical among males and females in each of the three populations.

#### Lactate dehydrogenasse (LDH)

Two bands were present at relative mobilities of 0.43 and 0.30. These were apparently monomorphic and of identical mobilities in all three populations of N. virescens from India, Sri Lanka and the Philippines.

and 0.41, in all three nonulations

Malate dehydrogenase (MDH) MDH showed two bands with very similar mobilities and no apparent variation among the individuals. The bands were of identical mobilities, 0.45

# Malic enzyme (ME)

Two slow moving bands were seen at relative mobilities of 0.20 and 0.16 Both were monomorphic with identical relative mobilities in all three N. virescens populations.

# Phospho2lucomutase (PGM)

PGM manifested two bands, but with very close mobilities and no variation among the individuals in the populations, Hence PGM could be considered to represent a single locus. The bands were of identical mobilities 0.24 and 0.19 in all three nonulations

#### Discussion

Of the eight enzyme systems studied, all except EST possessed bands with virtually identical relative mobilities in the three populations of N. virescens. The bands of the other seven enzymes appear to be homologous across the populations from India. Sri Lanka and the Philippines.

Est-1 had bands with virtually identical mobilities across the populations. Both Est-2 and Est-3, though postessing identical relative mobilities in India and Sr1 Lanka populations, differed from the Philippines propulation. Thus Est-2 and Est-3 gave diagnostic differences which are clear enough to allow separation of the Philippines population from the other two populations,

24

Of the total of 14 presumptive loci in the eight exayme systems, easy to Est-2 and Bix1, and differing relative mobilities between the two geographically different groups of India and Sri Lanka populations when conpared to the Philippieus population. This Intra-species genetic divergence between geographically distant populations of N virevers appears to be correlated with the partial reproductive lookation induced by differences in acoustical behavior. Yusof (1982) reported that pulse repetition frequencies of N virevers from Sri Lanka differed significantly from that of N. virevers population from Malaysia, which approximated the pulse repetition frequencies of N. virevers from the Philippines.

Claridge et al. (1984) also reported that pulse repetition frequency may be an important element endowing specificity to male calls in another homospieran, N. Ingens. Claridge et al. (1985) also reported that the difference between pulse repetition frequencies of N. Ingens were distinctive for populations from different geographic regions.

The more intense staining of Est.1 in the females indicates a higher degree of est-rate activity and hence Est-1 may be sax pelated in N. viezeux. Differential esterate staining intensity was reported between the males and females in S gatella fureignes Horwarh population from Sri Lanka on polycylamide gels by Rajaed. mr (1990) who most a more intense staining among males Sc-linkage in este-ase has also been reported in the caddally Themselois natural Milline Steiner et act. 1983.)

# Acknowled: ement

Grafeful thanks are due to the following of University of Wales, Curilly, United Kingdom » Professor M. F. Charliqg, Head, School of Pare & Rapfilled Biology, for facilities; Drs. J. Den Hollander, D. R. Lees and M. P. Wilnow for belight Comments and Mr. J. C. Morgan for retentical assistance by the control of the contr

Bellocher, S. H. (1979) Biochemical approaches to strain, race and species discontinuous pp. 137-144. In: Hoy, M. A. & Mc Keby, Jr., J. J. (Eds.), Genetics in relation to insure management. Rockefeller Foundation, New York.
Charles, M. F., Den Hollander, J. & Morgan, J. C. (1984) Specificity of

acoustic signals and mate choice in the brown planthopper Nileparrasia legent. Entimol. exp. & oppl. 35, 221-226. Chridge, M. F., Den Hollander, J. & Morgan, J. C. (1985) Variation in courtship signals and hybridization between geographically definable populations of the jick brown planthopper, Nileparante juepus (Sal.) Bid. J. Linn.,

Soc. 24, 35-49.

Ghauri, M. S. K. (1971) Revision of the genus Nephotetitix Matsumara (Homoptera: Cicadelloidea: Euscelidae) based on the type material. Bull. Entomol. Res. 60, 481-512.

lnoue, H. (1986) Biosystematic study of the genus Nephotettix occurring in Asia. Bull. Kyushu Natl. Agric. Expt. Stn. 24, 149-237.

Kalode, M. B. (1983) Leafhopper and Planthopper pests of rice in India, pp. 225-245, In. Proceedings of the 1st International Workshop of Leafhoppers and Planthoppers (Lukemortynicha) of Economic Importance, Commonwealth Institute of Entomology, London.

Khush, G. S. (1979) Genetics of breeding for resistance to the brown planthopper. In: Brown planthopper: threat to rice production in Asia. International Rice Research Institute, Los Banos, Philippines.

Ozaki, K. (1969) The resistance to organophosphorus insecticides of the green rice leafhopper, Nephotentix cinetice pt Uhlet and the small brown planthopper Landelphas striatellus Fallen. Rev. Plant Protection Res. 2, 1–15.

Powell, J. R., Tabachnik, W. J. & Arnold, J. (1980) Genetics and the origin of a vector population: Addes argypti, a case study. Science 208, 1385-87.

of a vector population: Aedes aegyptl, a case study. Science 208, 1385-87.

Rajendram, G. F. (1987) Esterase variation in Nephotettix nyropictus (Stal) population from Australia, pp. 303-309. In Proceedings of the 6th Aucoenorrhyncha

tion from Australia, pp. Meeting, Turin, Italy. Horvath population from Sti Lanka. Insect Sci. Applic. 11 (6) (In press).

Shaw, C. R. & Prasad, R. (1970) Starch gel electrophotesis of enzymes - a compilation of recipes. Biochem. Genet. 4, 297 - 320.

Steiner, W. W. M., Brigham, A. R. & Unzicker, J. D. (1983) Genetic studies of the caddisfly Triaenodes taraus Milne (Trichoptera: Leptoceridae). I Evidence for a sex-linked esterase. Biochem. Genet. 21, 1203 – 1207.

Yusof, U. M. (1982) Biological and taxonomic studies on some leafhopper puts of rice in southeast Asia. Ph. D. thesis. University of Wales, Cardiff.

Received for publication May 1990



# SOME OBSERVATIONS ON THE POWDERY MILDEW DISEASE OF SESAME (SESAMUM INDICUM L.)

# JEYADEVI KAILAYAPILLAI,1 A. SIVAPALAN 2 & K. THEIVENDIRARAJAH

(Department of Botany, University of Jarfna, Sri Lanka)

Vingnanam — Journal of Science 4: 27-38 (1989)

ABSTRACT: This study describes the morphology of powdery mildew fungus at various developmental stages in its life cycle and the symptoms produced on different parts of the sesame plant. Under natural conditions the infection first appeared as small tiny white superficial natches on the unner surface of mature leaf of a plant at the age of 40 days or more. Symptoms include surface lest necrosis, premature leaf fall, stunted growth of the plant at early stane vellowing and chiorosis of leaf at mature stage and browning of flower buds. Life cycle of powdery mildew was initiated by air-borne conidia, measured 25-36 \$\mu\_m\$ in length and 14-18 #m in breath. At 25-1°C and 100% R H. conidia started to perminate in about 4 hours and germination was maximum at about 24 hours after incubation. The penetration of the host tissue was completed within 28 hours. The second germtube was observed after about 30 hours and the third after about 36 hours of incubation. Secondary elongating hyphae, conidiophore initials and abstriction of conidiophore were observed 60, 144 and 155 hours after incubation respectively. The colony produced maximum amount of conidia on the ninth day after incubation. The colony remained productive even upto 12 days from the time of inoculation.

#### Introduction

One of the important diseases of seasme / Scientum Indicam L.) is the powdery mildew disease. The disease is caused by a fungus of which the contidial stage is Oddhim and belongs to Erysphaceae. The fungus responsible for seasme powdery mildew was named Oddim erysphoid by Rey (1963), Oddhim species by Secuevistrae and Appedrari (1964) and Levellule autors by Rangaswamy (1979). The fungus in the present study was named Oddim species following the identification at Commonwealth Mycological Institute, U. K.

Present address: Regional Office, Department of Agrorian Services, Puttur, Jaffna.
 Prisent address: 2/69 Noil Street, Greeneboough, Victoria, Australia.

The fungus is an obligate parasite which cannot be cultivated is no. Some of the marked characteristics of powdery mildew are suspensial physics mycelium, haustoria in the spidermal cells of the hosts, inxuriant development in dry weather, high water content of the air borne condia, diurnal perioders, with respect to several characters, compatible association with thirt hosts and vulnerability to control by fungicides (Yarwood, 1957). The white superfixing principles of the powdery mildew make them among the most easily recognized and best known fungi (Yarwood, 1959).

The powdery mildew disease is observed in almost all fields in Jaffar groin during January to July as essame is cultivated in the area during these particular months. The disease was observed in the glass-house throughout the year. The wide prevalence of this disease in Sri Lanka, particularly in the Jaffan district, and the severity of the disease at times have prompted this investigation.

# Materials and Methods

Observations were made on the morphology of powdery mildew at various developmental stages in its life cycle and the symptons produced on different parts of sessime plants. This study was carried out in field and in laboratory. Observations were taken in the fields of University of Juffas, Thriumelvely, and in Navali and Velbanai. For laboratory experiments, the level of the sessme plant of variety TMV<sub>3</sub> were used as the source of host tissue unless otherwise stated,

#### Source of inoculum

98

Sesame plants of variety TMV<sub>2</sub> grown in field and pots were allowed to develop natural infection and infections were continuously maintained on fresh potted plants throughout the experimental period

## Method of inoculation

The heavily infected leaves were gently shaken to remove old confision and were placed on filter paper diese in clockep plett dishes in all lluminated incubator. The following day the infected specimens were removed from the incubator and taped over a cardboard olysineer of 80 cm diameter and 2 in health follow which the healthy leaves (upper surface facing upwords) and most filter paper diese, with the montained leaf discipleness were placed dishes. The filter paper diese were kept were facilities that the face facing granted part of the contraction of the face of the contraction of the face of the

29

December 1989

Bedbatton

Sets of inoculated leaf discapleaves were incubated at 25°=1°C and

100°C RH, in an illuminated incubator, Inoculated potted plants were kept

a a glass-boase in a humid environment, where humidity was maintained

part 100% by placing buckets of water.

### Assessment

At the end of incubation, the inoculated leaf materials were taken out of the incubator or were brought from the glass-house and observed under the light microscope.

### Germination of conidia

The inoculated leaf materials were taken out of the incubator at two bour inearwals starting from the time of inoculation and were examined for grainstion of couldin. Collotage impressions of the inoculated surfaces were taken, staired with cotton blue in attorphenol, mounted in glycerine and observed under the light microscope.

The number of germinated and ungerminated costifis were counted with the holp of a tally counter. Germination counts were made form ten microscopic fields (x 100) selected at random on each of the ten leaf dissrabout 2500 contidia were observed for each treatment and a comition was considered to have germinated when the length of the germ time succeeds its breadth (Manners and Hossain, 1963). The mean percentage germination of ondition after different periods of incubation was acciousmed.

### Development of infection

Firstly the growth of germ tube was observed at two low show shows a starting from the time of insociation up to 24 hours. Inevented we are large were removed from the incubator at deficest time showns, assume superstions of the innociated surfaces were take and the innocessor, which will will noticology and the innocessor with the start of the shown and and the innocessor will be start in the start of the shown and the innocessor will be start of the shown and the shown an

## Production of conidiophore

Healthy leaves of sesume incoulated with positions with a session tapping method were incubated under optimal conditions in an incompanied

incubator. Samples were taken out at 24 hour intervals and were observed for the formation of conidiophore initials on the colonies. Camera lusted drawings of the colonies were made under light microscope after 144, 168

Apart from these observations, quantitative measurements on the number of conidiophores produced per colony in one sq cm area of the box tissue of all five replicates, after six, seven and eight days of incubation were determined.

#### Production of conidia

20

Potted plants of seature were raised from seeds under glass-long conditions. Five seedings of five leaf stage were incontated simultaneously to the tapping method and incobated at room temperature. After five days of incotation incodated leaves of the ages two and three were increased random from each plant and incobated at 25°C and 100% RH in an illuminant incubator for further seen days. Three one sq om cellotape impression were made from each of the five leaves and the number of frest codial profused per sq om area of the leaf on the 7th, 8th, 9th, 10th, 11th and 12th day after incudation was counted.

#### Experiments and Results

The disease was observed during February to June 1983, February to July 1984 and January to July 1985. In 1985 the disease was not observed during the month of June. The disease was observed under glass-house conditions throughout 1983 and 1984.

The disease usually appeared in the field in which the age of the same plants was more than ad Oasy. Under natural conditions the lower leaves developed infection first and from these the infection spread to the lower leaves and finally to the cities parts of the plant. When the temperature of the environment was high the infection appeared on the lower sufficient of the upper leaves and both surfaces of the lower leaves, about 10-40 drift first large parameter of infection the young leaves also developed infective first large parameter of young leaves also developed infection that the lower surface has on the upper sufficient the possible of young branches developed infection in a way similar to thors the proper parameter of young branches developed infection in a way similar to thors the proper parameter of young the property property of the property pr

#### Symptoms

Symptoms

Powdery mildew infection on sesame plant has been apparent from seedling stage up to mature plants. All parts of the sesame plant namely the

gaves, stem, flower bads and pods were found to be affected by this fungus, Leaves were found to be the mass ususceptible tissue to fungal attack. The first symptoms of the infection were the small timy white almost occurat superficial powdery patches on the host surface. These white putches increased in number and site and became puvdlery in nature. These powdery patches collected and eventually covered the entire leaf surface. The vibble white putches appeared a day before the appearance of the sprovinging structures. The affected portions of the leaves became dry, turned bown and later developed marginal necrois.

The entire leaf surface already brown and dry curl.d up and finally dropped off from the plant leaving only the stem. In some cases the heavily infected leaves before showing symptoms of necrosis became pale in colour and dropped off from the plant.

Sesame plants infected at an early stage of growth (about 20 days) showed reduced or stunted growth. On one coasion in a field where 20 day old plants developed infection, the cultivation was completely abundoned about three weeks after the initial appearance of infection. The infection but bads became brown, withered and failed to open and develop into podaring the period of severe disease incidence the pol infection above was observed. Small pods were more sunceptible than the matter pods. Due to the pol infection infected pods became pale green in colour and were constituted with the powdery mass of the fingus. These pods produced fewer, thinner and smaller seeds than the normal ones.

Infected young stem also was observed in heavily infected setame fields.

Affected stems were covered with powdery mildew fungus. In addition they became slender and lost their rigidity. In such stems only few flower buds were produced.

Life cycle of the fungus

Germination, infection and sproulation follow each other in the life cycle of powdery mildew.

#### Germination

Life cycle of the powdery mildew is initiated by airborne condition which are the asexual reproductive spives of the fungus. The condisploner were diliptical, smooth walled, colouriess and vaccoulsted when fresh. The Publice condition measured 25 to 36 Pm in length and 14 to 18 Pm in width. When the conditions were favourable germination of the fungus was observed as direct germination of conidia.

39

The mean percentage germination of conidia after different periods, of incubition was calculated. The condidia started to germinate after about four bours and produced a single thick germitube from one corner of the coniding. The mean percentage germination of conidia interested with increase in period of incubation and attained maximum level of 70% around 24 hours after incubation (Fig. 1).




Fig 1 Germination of conidia of Oidium s ecies on leaf discs of sesante after different periods of incub tion, at 25°C and 100% RH.

33

## Infection

Firstly the growth of the germinbe was observed at two hour intervals from the time of incondution up to 24 hours. From the data the mean length of germinube after different periods of incubation was determined (Fig. 2). The observation revealed that the coulding germinated after about four hours of incubation and the germinube continued to grow and attained in maximum length at about 10 hours after incubation. Therefore the mean length of germinute constitued more to the continued up to 24 hours. The about 25 hours after incubation.



Period of incubation (Hrs)

Fig. 2 Growth of germ tube of Oidlam species on leaf discs of sesame, at 2 hour intervals upto 24 hours of incubation at 25°C and 100% RH.

Since the germinated conidia appeared to have formed additional germinate after 30 hours of incubation, it was necessary to continue the experiment further to trace the germinating conidia. Another experiment was set up in which the measurements on the hyphal growth produce at 24 hour intervals. From these measurements the mean total length of hyphae of each intervals. From these measurements the mean total length of hyphae of each

germinated condition at 24 hour intervals was determined. There values at expressed in the form of a graph (Fig. 3). Mean total length of hyphae increased with increase in period of incubation. Germinating conditions byphae at about 30 hours after incubation, the proposed of incubation of the proposed of the proposed




Fig. 3 Growth of hyphae of Oidium species on leaves of sesame at 24 hour intervals at 25°C and 100% RH.

## Sporulation

34

Camera lucida drawings of the colonies were made under light microscope. Conidophore initials appeared 144 hours after inoculation and the typically unbranched and septate condiciophores were formed after 160 hours of incoher tion. The mature one celled conidia were ready for release after about 100 hours of incoherance after a south 100 hours of incoherance after a south 100 hours after incoherance after incohera



Camera lucida drawings (×100) of a 198 hour colony of Oidium species on sesame leaf at 25°C and 100% RH.



Apart from these measurements, the number of condicipones produced per colony in one sq can area of host tissue-after its, seven and eight of incubation were determined. The results in Table 1 revealed that the rate of production of condisplones increased with increase in period of inoubation. The mature condica were ready for release by about seven days after inoculation and the colony remained productive for a further five days.

Table 1: The number of conidiophores produced after different periods of incubation 25°C and 100% R H.

Hours after incubation	Number of conidiophores Produced / colony
144	strance to the pare mans off
192	et dich atr 780 heurenbare bar all id

The values denoted by different letters are statistically different.

The number of fresh conidia produced per sq cm area of the last on the 7th, 8th, 7th, 10th, 11th and 12th day after inoculation was counsel. The results in Fig. 5 indicated that the rate of fresh condili produced intreased up to nime days and then there was a gradual decrease in the condition production up to 12 days. Therefore the establishment of the fragges from the time of inoculation to spore release on leaves was found to take nimedays.




Fig. 5 The production of fresh conidia of Oidium species on leaves of sesam at 25°C and 100% RH after different periods of incubation.

## Spore release

The conidia were released when the host leaf and the atmosphere were relatively dry, that is during the day time. Detached conidia were taken away by the wind and deposited over the fresh host leaves on which they germinated seain.

Sexual reproductive structures of the fungus such as perithecia were not observed during the three years of study.

#### Discussion

The present studies have established that the consuirs organize of the powdery milities disease of seasure in Jaffas is the Outen 8... Though the disease was widespread throughout the year, the seventy of the disease was widespread throughout the year, the seventy of the disease was found to be high during the months of September to February. The relatively lower temperature, high relative humidity and reduced sunlight experienced during these months seem to favour the development and good the properties of the control of the

The fungus was able to establish itself on all parts of the plant but it generally starts on the more matter lower leaves of the stem and on the upper surface. It then quickly append to both surfaces of the lower leaves and to other leaves, main stem, beanches, flower bods and even mature pols. Similar observations, affecting all parts of the plant on a number of pols. Similar observations, affecting all parts of the plant on a number of Arraycod. 1973 and Wesslers (1964).

This fungas has a relatively short life cycle of about 12 days. The infection starts with the germination of the condition on the plant surface. The condition begins to germinate after four hours and produce white byphal masses visible to the niked eye in five days. The specilism starts production condition from the 7th day until the 12th day with a maximum production on the 9th day. The condition were released when the host leaf and atmosphere was the condition of the start of the

It was also observed that beavy raises wiped out the infection but on after the raises the infection on appeared only on the lower surface. Mild raises for short period on the other hand lead to luxurisat development of providery mildew on host surfaces. Greater spacing between pileats surface over crowding and shade coupled with see frequently of the discussion of the coupled with see frequently of the discussion of the coupled with section of the coupled with section of the coupled with the provider of the coupled with the co

#### References

Cook, H. T. (1931) Powdery mildew disease of snap beans. Virginia Truck Agr., Exp. Sta. Bull. 74, 931 - 940.

Kailayapillai, J. (1987) Studies on the Powdery mildew disease of seame
(Sesamum indicum L.). M. Phil. Thesis, University of Jaffina.

Vol. 4. Nos. 1 & 2 Manners. J. G. & Hossain, S. M. M. (1963) Effects of temperature and humidity on conidial germination in Erysiphe graminis. Transactions of

Vingnanam - Journal of Science

38

the British Mycological Society 46 (2), 225-234.

Miller, P. A. (1939) An important Powdery mildew disease of the coast live

oak, 6th West Shade Tree Conf. Proc. 1-5.

Rangaswamy, G. (1979) Diseases of oil seeds pp. 339-342. ln; Diseases of

crop plants in India, Rangaswamy, G. (Ed.) Perentice - Hall of India Private Limited. New Delhi.

Roy, A. K. (1965). Occurrence of Powdery mildew caused by Oidium - eryslohoides

Plant protection Bulletin F. A. O. 13, 42. Senewiratne, S. T. and Appadurai, R. R. (1966) Gingelly pp. 361-366, In:

Field Crops of Ceylon, Lake House Investments Limited, Colombo.

Wheeler, B. E. J. (1969) Powdery mildews, pp. 85 - 107, (In: An Introduction to Plant - Diseases. B. E. J. Wheeler (Ed.) The English Language Book Society and John Wiley & Sons Ltd., U. K.

Yarwood, C, E. (1957) Powdery mildews. The Botanical Review 23 (4), 235 - 300,

Received for publication May 1989

இதம் 4



1-ம், 2 ம் இவளியீடு

விஞ்ஞானம் விஞ்ஞான சஞ்நிரை

'கிஞ்ஞானம்' இதழ் ் இன் வெளியிடு எயக்கு அப்பாற்பட்ட குழ்நினவகளினாக் தாமதுமாகுவிட்டது. இவற்றில் மிகவும் ஆக்கரமான நிகழ்ச்சி முன்னால் பிரதம ஆகிர் யர் S. கந்தையா அவர்களின் நிடர் மடைறவாகும். இவர் 'கிஞ்ஞானம்' உந்பத்தியான காயத்தியிருந்த இப்பணியேன. ஆற்றினார்.

- இதழ் 4 இம் பௌதிகத்தில் முதனாவது கட்டுரையை வெளிபிடுகிறோம்: K. அந்தசாயி: ''உனோகங்களில் ஐதரசன் பரவல்''.
- இரு கட்டுரைகள் இப்பிரதேசத்தின் பலிர்களின் தோய்களைக் குறிக்கும்: K. V. S. கத்திரசன், பத்மாசனி வரதாரஜாவின் கட்டுமை பப்பாகிலின் நிதிராவுவு தொகும்பும், தெறுபிதனி கணைபபின்னை, K. கிவாமண், K. தெறிவேதிரராஜாவின் கட்டுரை என்னக் காவரத்தின் சாம்பிடிழ்ந்தனை நிருமையும் தெரிக்கும்.
- G. F. இரசிஞத்தும், நீர்மன R. ஆன்ரனி, நோவின் நகவிலம், ஆபெருடன் யாற்ப்பாணக்கு டாரட்டின் துனம், பற்றிய சூமாயது ஆல்வை ஆர்கித்துள்ளர். G. F. இரசிஞத்துக்கில் மற்றாற் உடற்றை இந்தும், இவக்கை, பிலிமையின் நாடு கவிலுள்ள ஒரு பச்சா இணைத்ததின் குடித் தெராவகவிடமெறுள்ள புலிம்வி சீதியான முறியனரியின் வெறுர்களைப்பற்றிக் தெளிகிகின்றது.

aved 1991



g Diffe

Vol. 4, Nos. 1 & 2 40 Vinguanam - Journal of Science December 1989

Vingnanam J. Sci. 4 (1989)

இவ்விதம் கட்டுரைகளின் சுருக்கங்கள்

உலோகங்களின் ஐதரசன் பாயம் அளியர் : K. adagoul

(Qபளடுகளியல் அறை, யாழ்ப்பாண பல்கலைக்கழகம், இலங்கை.)

Vingnanam J. Sci. 4:1-7. காக்கம்: உளோகங்களில் ஐதரசன் பாயத்தின் பரவனுக்கான சமன்பாட்டை பேஸலகங்க

தகைப்புக்குட்பட்ட நிண்மத்தில் அசையும் கரை பொருள் பாயத்துக்குரிய சென்மில். லிவின் முறை இயல் உபயோகப்படுத்தப்பட்டது. இச் சமன்பாடுகள் ஆரவில் குழ வினரால் அறிகிக்கப்பட்ட பர்சோதனைத் தரவுகளுடன் ஒப்பிடப்பட்டன. இசைவுற தன்மை இருப்பியாகவுள்ளன.

பப்பாலி சித்திரவடிவு நோயினால் தொற்றுகையுற்ற பப்பாலி (CASICA PAPATA L.) இலைகளில் இருக்கும் சுயாதீன அமினோ அமிலங்களின் ஆராய்வு

அசிரியர்கள் : R. V. S. சுந்தரேசன், பத்சைனி வரதாஜா (காவரவியல் துறை, யாழ்ப்பாணப் பல்கலைக்கழஈம், இலங்கை)

Vingnanam J, Sc. 4:8-13.

எருக்கம் :

Carica papaya L, C. V. Coimbatore 2, C. P. 124, Local, Solo Hawaii, Unknown Indian ஆகிய குல வகைகளின் ஆரோக்கியமான இலைகளினதும் பப்பரி இவைச் சித்திர வடிவு நோயினால் தொற்றுகையுற்ற இவைகளிரதும் தெடுப்பினை அயன் மாற்றிடு திறப்பகுப்பியல் செய்தபின்னர், இரு பரிமான தாழ் நிறப் பகுப்பியல் முறையினால் சயாதின அமினோ அமிலங்களின் இருக்கைபற் நித் துணியப்பட்டது. Alanine, « - amino butyric acid, arginine, aspartic acid, cysteic acid, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenyl alanine, serine, threonine, tyrosine, valine ஆகிய அமினோ அமின களின் இருக்கை ஆரோக்கியமான இவைபினதும், தொற்றுகையுற்ற இடையினதும் பி∮த் தெடுப்பிலும் காணப்பட்ட போறிலும் அவற்றின் செறிவு வெல்வேற குளவகைக்குக் இடையி**க்** வேறுபட்டது. இல ஆரோக்கியமான குலவகையிலும், தொற்றுகையுற்ற கு<sup>ல</sup> வகையிலும் அமீனோ அயிலங்கள் இக்காமையும், செறிவு மாறுபடுதலையும் காரணம்

கூறி விளக்குவதற்கான முயற்கென் மேற்கொள்ளப்பட்டவ.

இவ்விதழ் கட்டுரைகளின் சுருத்கங்கள் மார்கழி 1989 <sub>யாழ்</sub>ப்பாணக் குடாநாட்டின் வீட்டுப்புற நுளம்பு இனங்களின்

41

<u>ு</u>யிரியல் அசியர்: G F. Agrifo ந்திரம், நிர்மலா R. அன்றவி, நேடினி நாகலிங்கம்

(வீலங்கியல் துறை, யாழ்ப்பாண பல்கலைக்கழகம், இலங்கை) Vingnanam J. Sci, 4:14-19.

agtsù:

யாழ்ப்பாணக் குடாநாட்டின் மூன்று இன வீட்டுப்புற நுளம்புகளின் உயிரியல் ஆராவப்பட்டது. Aedes aegypti Linnaeus, Armigeres durhami Edwards, Culex quinquefosciatus Say. A. aegypti மின் பக்வேறு நிலைகளின் சராசரி விருத்டுக் கால மானது: முட்டை 1.69 நாட்கள், குடம்பி 4.5 நாட்கள், கண்டுப்புமு 6.7 நாட்கள். போத்தவிருத்திக் காவம் 12.9 நாட்கள். A. durhami வின் விருத்தி நிவைகளின் சரா சரிக் காலம்: முட்டை 3.12 நாட்கள், ஆடம்பி 6.84 நாட்கள், கண்டுப்புஷ 4.04 நாட்கள். மொத்த விருத்டுக்காலம் 14.0 நாட்கள். C. quinquefasciatus இன் இளமைப் பருவங்களின் சராசரிக் காலப்: முட்டை 5.6 நாட்கள், குடம்பி 5.8 நாட்கள், உண்குப் புழு 2.9 நாட்கள். மொத்த விருத் தெக்காலம் 14.3 நாட்கள். மூன்று இனங்களினதும் திருத்திக்காலம் வெப்பநிலையில் கங்கியிருப்பகாகத் காணப்படுகிறது.

இந்தியா, இலங்கை, பிலிப்பைனின் Neph-tettix virescens (Homoptera: CicadelliJae) குடித் தொகைகளின் நொதியங்களின் மின் பகுப்பு ஆய்வு அசெயர்

G. F. @g 1 700 i Strik (விலங்கியல் துறை, யாழ்ப்பாண பல்கலைக் பூசம், இலங்கை.)

**ஈாக்கம்**:

Vingnanam J. Sci. 4: 20-26. எட்டு கரையக்கூடிய தொடுயங்களின் பட்டிகை மாடுரிகன், என்ரேசு (Est), குகுக்கோக 6 பொஸ்பேற் மறுதாகணேக (G-6-PHD), எக்ஸ்சோகைணேக (HK), ஐசோகத்திரேற்று முறுதரசனேக் (IDH), மிலற் மறுதரசனேக் (MDH), லகிரேற் ம.ஐதரசனேசு (LDH) மலிக் தொடுயம் (ME), போன்போஜ சூக்கோமியூட்டேசு (PGM) ஆகியன, இந்தியா, இவங்கை, பினிப்பைன் நிவுகளிலிருந்து சேகரிக்கப்பட்ட Nephoterix virescens (Distant) குடித்தொகைகள், ஐக்கிய இராச்சியத்டுலுள்ள, காடிவ்ப், வேல்ஸ் டல்கலைக்கழகத்தில், கண்ணாடி – வளர்ப்பு இல்லங்களில், TNI தெல்லினத்தில், பராமரிக் கப்பட்ட வளர்ப்புகளில் ஆராயப்பட்டன. இரு இலிங்க நிறைவுடலி பூச்சிகளிலும், எட்டு நொடுவங்களுக்கான மாப்பொருள் ஜெல் மின்றுண்பகுப்பையும் (electrophoresis) ெற்ற சாயமுட்டவையும் பாவித்த இல்லாய்வு மேற்கொள்ளப்பட்டது. ஏழு தொடுயங் salar பட்டிகைகள், மூன்று குழு பதெ:கைகளிலும், எண்ணிக்கையிலும், சார் அசைவிலும் இணத்தமடைத்திருத்தன.

எஸ்ரெகில், ஒன்ற வையக்களில் தொழிலாடு என்ன என்ற அக்கொண்டிய கும் அமதாவிக்கப்பட்டது. இருட்டிகைகளில் என் அவவு, கிலிமணன் நடிக்கிலை பெறப்பட்ட அதிதொகையில் இந்தியா, இவக்கைகின் 87 weems ஒவிருந்த போறப்படுகுத்தது. அத்தடன் பிலிமைன் அதுகிறாகவை இவக்கள்ற மேறுந்துத் இது பயன்படுத்தியடனால். என்றோர் 1 பட்டிகைகள் இவிக்கத்துடன் செருபிறும், மனவாக, பெண் பூசிகளில் அவை அதிவனின் சுண்மபாகவும் நென்றின். இப்பெற பேறுகள் முன்ன அதுகாவிக்கப்பட்ட இவக்கை கிலிமணம் இறைக்கிறும் பெறப்பட்ட N. weems இன் அமிலியல் நடத்தையுக்கும். இரு குடித்தொகைவுக்கியடல்லை பதுதி இவல்போத்துக் அகியில் நடத்தையுகுற்க்

எள்ளு தாவரத்தில் தோற்று**வி**க்கப்படும் சாம்பல்பூஞ்சண நோயின் சில அவதானங்கள்.

ஜெயதேவி கைலாயபிள்ளன. A. வெயாலன், K. தெய்வேற்றிரசாஜா. (தாவரலியற்றுறை, யாற்ப்பாண பலிகலைக்கழகம், இலங்கை) Vingnanam J. Sci. 4: 27—38,

கருக்கம் :

அதிர்கள் :

என்னு தாவரத்டும் சாப்பம் பூஞ்சணத்டுன் வாழ்க்கைவட்டத்டும் வெவ்வே**ற** விருத்ர நிலைகளின் உருவனியலையும், என்னு தாவரத்றின் கெவ்வேறு பாக கணிக் ஏற்படுத்தும் நோயறிதறிகளையும் பற்றிய படிப்பு மேற்கொள்வப்பட்டது. இயற்கை நிபந்தனைகளில் தொற்றுதலானது 40 வயது அல்லது அதற்கு மேற்பட்ட தாவரத் தின் முதிர்ந்த இலையின் மேற்பரப்பில் சிறிய துண்ணிய வெள்ளைப்படையாக முத லில் தோன்றியது. இலைப்பரப்பு இநத்தழிதல் முன்முடுர்ந்த இலைவீட்சி, ஆரம்படிலை யில் தாவரத்தின் குறன் வளர்ச்சி, மஞ்சனாதல். முடுர்ந்த நிலையில் இவையின் பச்சை அழிதல், பூ அரும்புகளின் மண்ணிறமாதல் ஆகியன தோயறிகுறிகளுள் அடங்கும் இச் சாம்பல் பூஞ்சணத்தின் வாழ்க்கை வட்டமாதை 25 — 35 ∰ m நீளமும் 14 − 18 ∯m அண்டும் உடைய வளியிலுள்ள தூனிய வித்துக்களால் தோற்றுவிக்கப்பட்டன தூனிய வித்கொளைது 25±1°C மிலும், 100% சாரீரப் பதனினும் - மணித்தியாலத்டும் முளைக்க ஆரம்பித்தன. 24 மணித்தியால அடைகாத்தவின்பின் முனைத்தல் அடு உயர்வாக இருச் தது. இந்தோயாக்கியானது விருந்து வழங்கியில் 28 மணித்தியாவத்தில் ஊடு நமிபது. இரண்டாவது மூலவுயிர்க் குழாயானது 30 மணித் யாலத் `ன்பின்பும் மூன்றாவது 🕬 வுயிர்க்குழாயானது 36 மணித்தியால அடை காத்தலின் பின்பும். அவதானிக்கப்பட்டது. துணையான தீளும் பூஞ்சண இழை, தூனியந்தாங்கி முதல்கள், தூனியந்தாங்கியின் கடிங் கியனுதல் ஆகியவ முடையே 64, 144, 156 மணித்தியால அடைகாத்தலின்பின் கூற தானிக்கப்பட்டன. சமுதாயமானது 9 நான் அடைகாத்தலின்பின், அடுகூடிய துனிபவித் இகளைத் தோற்றுவித்தது சமுதாயமானது உட்புகுத்துவிலிருந்து 12 நாட்களின் பின்பும் உற்பத்தியாக்கும் இறனைக் கொண்டிருந்தது.

## VINGNANAM - JOURNAL OF SCIENCE

## Instructions to Contributors

#### GENERAL

Submission of a manuscript to the editor involves the assurance that it is original and that no similar paper, other than an abstract, has been or will be submitted for publication elsewhere without the consent of the Editorial Board.

Research papers from original investigations and reviews which are critical evaluations of existing knowledge in a specialised field are accepted for editorial consideration. Short communications of sufficient importance to merit publication in advance of a full paper are also accepted by the journal.

The language of publication is English. A translation of the abstract in swabs has should be submitted with the manuscript. Every paper will be referred to at least one referree familiar with the field of research covered by the paper, who will be nominated by the Editorial Board. Papers are edited to increase clarity and case of communication.

## PREPARATION OF MANUSCRIPTS

No restriction is laid on the length of the manuscript, provided it is written clearly and concisely. All unnecessary descriptions, figures and tables must be climinated. The innersonal form should be used in the text.

The style of setting out, sub-division of text and lay out of tables in the manuscripts should in general be organised in the form adopted in this issue.

Manuscripts should be submitted in triplicate including the original typewritten copy, typed throughout in double spacing on one side of the paper only. Adequate margins (4 cm) should be left with liberal spacing at the top and bottom of each page. The typescript should be free of corrections.

Each page of the manuscript should be numbered in the upper right hand corns of the page. The last page should contain (a) a note as to the number of numscript pages, figures and ables, (b) proposed running title of not more than four words and (e) the name and mailing address of the person to whom the proofs thould be sent?

#### ILLUSTRATIONS

All illustrations are considered as figures and each graph, drawing or photograph should be numered in sequence with Arabic un uterals. Authors must submit the original and two duplicates of each figure. Figures should be planned to fit the proportions of the printsd page. The maximum space available on a page is 140 x 190 mm. Figures must be drawn in Iralian ink on plain white paper or board trades paper, not larger than twice the linear dimensions desired. Drawings should be paper, not larger than twice the linear dimensions desired. Drawings should be lettered with a lettering successful the paper of the larger should be been paper of the larger than the letters and numerate plain and the larger than the larger tha

Each figure should carry a legend so written that the general meaning of each illustration can be understood without reference to the text. The amount of lettering on drawing should be reduced as far as possible by transferring it to the legend, Figure legends should be typed on a separate sheet and placed at the end of the manuscript.

Graphs should be plotted on white or bluelined graph paper or tracing cloth; grid lines that are to be shown in the engraving should be inked in black. The caption of each axis should be lettered parallel to it saxis. Each figure should be identified in the margin with figure number. The preferred position of all ulturarious should be indicated in penal in the manuscript.

#### PHOTOGRAPHS

Good glossy prints with sharp contrasts between black and white areas should accompany the manuscripts. The size should be such that when the print is reduced to the normal size for reproduction (140 × 190 mm maximum) the detail is still clear. Magainfeation should be indicated with a scale line on the photograph. Fisure number should be siren on the back of each photograph.

## TABLES

Tables should not repeat data which are available elsewhere in the pator. Each table should be typed on a search schw-tift doe regard for the prototions of the printed page. They should be numbered consecutively with Arable mannests. Tables der matter should be cately see out and the number of colorium control of the colorium of the colorium control of the colorium colorium control of the colorium coloriu

## REFERENCES

All references to publications made in the text should be presented in a list of references following after the text of the manuscripts. The manuscripts should be carefully checked to ensure that the spelling of authors names and dates are exactly the same in the text as in the reference list. In the text refer to the author's name (without initial) and year of publication, followed, if necessary, by a short reference to appropriate pages. E.g. "Since Peterson (1967) has shown that, ..." "This is in agreement with results obtained later (Kramer, 1969, pp. 12-16)". If reference is made in the text to publications written by more than two authors. the name of the first author should be used, followed by "et al.". This indication, however, should never be used in the list of references, where the name of authors and co-authors should be mentioned instead. The list of references should he arranged alphabetically on authors' names, and chronologically per author. If an author's name in the list is also mentioned with co-authors, the following order should be used. Publications of the single author, arranged according to public ation dates, publications of the same author with one co-author and publications of the author with more than one co-author. Use the following system:

### IOURNAL. Selman, I. W. & Kulasegaram, S. (1967) Development of the stem tuber in

Kohlrabi. Journal of Experimental Botany, 18, 471 - 490, Journal name should not be abbreviated BOOK

## Slatyer, R. O. (1967) Plant Water Relationships Academic Press, London, 366 pp.

SECTIONS OF BOOK

Skoog, F. & Miller, C.O. (1957) Chemical regulation of growth and lorgan formation in plant tissue cultured in Vitro. pp. 118-131. In Symposia for the Society of Experimental Biology XI. The Biological action of growth substances University Press, Cambridge,

In referring to a personal communication the two words are followed by the Year, e.g. (J. McNary, personal communication, 1968).

## CROSS\_REFERENCES

The cross reference cannot be finally inserted until the page proof is available. Type them: "see page OOO". In the margin pencil the page number of the cross reference in the manuscript.

#### FORMULAE

Formulae should be typewritten, if possible, leaving ample space around the formulae, Subscripts and superscripts should be set off clearly. Greek letters and other non-Latin or handwritten symbols should be explained in the margin when they are first used. Take special care to clearly show the difference between zero (O) and the letter O, and between one (I) and the letter I. Give the meaning of all symbols immediately after the equation in which they are first used. For simple fractions use the solidus (/) instead of a horizontal line,

e.g.: 
$$1_p/2_m$$
 rather than  $\frac{1_p}{2_m}$ .

Parentheses and square brackets are preferred in formulae. Accoludes should be used only when they are absolutely necessary. Equations should be referred in frequency and the mobile referred to the property of the property

the symbols, e.g. 18O. Use of superscripts added to superscripts and subscripts added to subscripts should be avoided, if possible.

# SHORT COMMUNICATION

The journal may include a limited number of short communications. Another should submit short communications only when they believe that rapid publication of their result is of the utmost importance. A short communication must not exceed 1200 words, should be complete in its own right and suitable for citation.

#### PROOFS

Corrected galley proofs must be returned to the editor without dealy, Faulure to do to will result in delay in publication. Correction of pools by authors must be restricted to the printer's and similar errors. They should be marked in pencil. Any modification of the original text is to be avoided. Rev ponsibility for correcting proofs rests entirely on the authors though editorial sestiance will be provided.

#### REPRINTS

Twenty reprints will be supplied free of charge for each article. Additional toprints can be ordered on the reprint order form which will accompany the proof.



## VINGNANAM

December 1988

## JOURNAL OF SCIENCE

Numbers 1 & 2

38

42

CONTENTS	
Laboratory Studies on the effects of Environmental conditions and Fungicides on the Development of Alternaria alternata (Fr.)Keissler, NIRANJANI RAMANATHAN and A. SIVAPALAN.	i
Some Field Observations on the Epidemiology of Downy Mildew Disease of Grape Vine. NIRANJANI RAMANATHAN	10
Effect of Volatile Substances from Azadirachia indica (Neom) seeds on the Reproductive Biology of Tribolium castaneum (Colcoptera : Tenebrionidae). PATHMINI MAHESWARAN and V. K. GANESALINGAM	20
Isolation of Intra-Cellular Symbiotes from the Brown Planthopper Niiaparvata lingens (Homoptera: Delphacidae) occurring in Sri Lanka, G. F. RAJENDRAM and ANNETTE SELVADURAI	27
Observations on the Embryonic Development of Nephotestic virescens (Homoptera: Cicadellidae). G. F. RAJENDRAM and RAJINI RAJESWARAN	31

Instructions to Contributors

Abstracts in Tamil

Volume 3

Effect of Horizontal Centrifugation on Reproductive Biology of Callosobruchus
maculatus (Fabricius) (Coleoptera: Bruchidae). SAKUNTHALADEVI
AMBIKAIPAHAN and V. K. GANESALINGAM

Heat Stress in Sri Lanka - A Human Climatic Approach.

MANICKAM PUVANESVARAN

#### VINGNANAM

### IOURNAL OF SCIENCE

Volume 4

December 1989

Numbers 1 & 2

#### CONTENTS

Diffusion of Hydrogen in Metals.

A Study of the Free Amino Acids present in Leaves of Papaya (Carlea papaya L.) varieties infected with Papaya Mosaic Disease.

R. V. S. SUNDARESAN and PATHMASANY VARAIHARAJAH

Biology of Peridomestic Mosquito Species of Jaffna Peninsula.

G. F. RAJENDRAM, NIRMALA R. ANTONY and
ROHINI NAGALINGAM

Starch Gel Electrophoresis of Enzymes of Nephotettis Virescens (Homoptera: Cicadellidae) populations from India, Sri Lanka and the Philippines. G. F. RAJENDRAM

Some Observations on the Powdery Mildew Disease of Sesame
(Sesamum indicum L.). JEYADEVI KAILAYAPILLAI,
A. SIVAPALAN and K. THEIVENDIRARAJAH

Abstracts in Tamil

Instructions to Contributors